PART - II - FINAL

BRANCH - MATHEMATICS

Paper: I: FINITE MATHEMATICS AND GALOIS THEORY

(Revised Regulations from 2010-2011)

Max. Marks: 20

SECTION-A

Answer any four quesitons. Each question carries 5 marks.

- 1. Show that s(r,n) = s(r-1,n-1) + ns(r-1,n) where $r,n \in N$ with $r \ge n$.
- 2. Find the coefficient of X^R $R \ge 13$ in the expansion of $(x^3 + x^4 +)^6$.
- 3. Prove that the degree sum of all vertices in any graph is even.
- 4. Prove that a graph and its compliment cannot both be disconnected.
- 5. If $f(x), g(x) \in Z[x]$ are primitive polynomials, then prove that their product f(x)g(x) is also primitive.
- 6. Let $F=\mathbb{Z}/2$. Then show that the splitting field $x^3 + x^2 + 1 \in F[x]$ is a finite field with eight elements.
- 7. If $f(x) \in F(x)$ has r distinct roots in its splitting field E over F, then prove that the Galois group G(E/F) of f(x) is a subgroup of the symmetric group S_r .
- 8. Show that the Galois group of $x^4 + x^2 + 1$ is the same that of $x^6 1$ and is of order 2.

SECTION-B

Answer One questions from each unit. Each question carries 15 marks.

UNIT-I

9. a) Find the solution to $Y_{n+2} - 4Y_{n+1} + 4Y_n = 3.2^n$

b) How many ways can 25 people be assigned to 3 different rows with at least one person in each row?

(OR)

- 10. a) Prove that the total number of k-permutations of a set A of n elements is given by n(n-1)(n-2).....(n-k+1).
 - b) Let $A_i = \{ \Pi \in S_n | \Pi(i) \in X_i \}$ be the bad permutation for i. Use principle of inclusion and exclusion, find the formula for $p(X_1, \dots, X_n)$.

UNIT - II

- 11. a) Prove that there are eleven nonisomorphic simple graphs on four vertices.
 - b) Prove that a graph is bipartite if an only if it contains no odd cycles.

(OR)

- 12. a) Let M be the incidence matrix and A be the adjacency matrix of a graph G. Then prove that every column sum of M is 2
 - b) Prove that in a tree any two vertices are connected by Unique path.

UNIT - III

- 13. a) State and prove Gauss Lemma.
 - b) Let p(x) be an irreducible polynomial in F[x]. Then prove that there exists an extension E of F in which p(x) has a root.

(OR)

- 14. a) Let E be an extension of F. If K is the subset of E consisting of all the elements that are algebraic over F, then prove that K is a subfield of E and an algebraic extension of F.
 - b) Let $f(x) \in F(x)$ be a polynomial of degree ≥ 1 wit α as a root. Then prove that α is a multiple root if and only if $f(\alpha) = 0$.

UNIT - IV

15. Let H be a finite subgroup of the group of automorphism of a field E. Then prove that $[E:E_H]=[H]$.

(OR)

16. Prove that every polynomial $f[x] \in C[x]$ factors into linear factors

PART II - Final

BRANCH: MATHEMATICS

Paper - II: TOPOLOGY AND FUNCTIONAL ANALYSIS

(Revised Regulation from 2010-2011)

Max. Marks: 20

SECTION - A

Answer any Four questions. Each questions carries 5 marks.

- 1. Define a topological space. Let J_1 and J_2 be two topologies on a non empty set X. Then show that $J_1 \cap J_2$ is also a topology on X.
- 2. Prove that every sequentially compact metric space is compact.
- 3. Prove that in a Hausdorff space, any point and disjoint compact subspace can be separated by open sets, in the sense that they have disjoint neighborhoods.
- **4.** Prove that the space \mathbb{R}^n and \mathbb{C}^n are connected.
- 5. If M is a closed linear subspace of a normed linear space N and x_0 is a vector not in M, then prove that there exists a functional f_0 in N* such that $f_0(M) = 0$ and $f_0(x_0) \neq 0$.
- 6. If P is a projection on a Banach space B, and if M and N are its range and null space then prove that M and N are closed linear subspaces of B such that $B = M \oplus N$.
- 7. Prove that a closed convex subset C of a Hilbert space H contains a unique vector of smallest norm.
- 8. Prove that an operator T on a Hilbert space H is self-adjoint if and only if (Tx, x) is real for all x.

SECTION - B

Answer one question from each unit. Each question carries 15 marks.

UNIT-I

- 9. a. Let X be a topological space and A a subset of X. Let D(A) be the derived set of A. Then prove that
 - i. $\overline{A} = A \cup D(A)$; and
 - ii. A is closed $\Leftrightarrow A \supset D(A)$.

236-14

(1)

[P.T.O.]

b. If f and g are continuous real or complex functions defined on a topological space X, then prove that f + g, αf , fg are also continuous.

(OR)

- 10. a. State and prove the generalized Heine Borel theorem.
 - b. Prove that a closed subspace of a complete metric space is compact if and only if it is totally bounded.

UNIT-II

- 11. a. Prove that every compact Hausdorff space is normal.
 - b. State and prove Tietze extension theorem.

(OR)

- 12. a. Prove that a subspace of the real line \mathbb{R} is connected if and only if it is an interval.
 - b. Let X be a compact Hausdorff space. Then prove that X is totally disconnected if and only if it has an open base whose sets are also closed.

UNIT-III

- 13. a. Let N and N' be normed linear spaces and T a linear transformation of N into N'. Then prove that the following conditions on T are all equivalent to one another:
 - i. T is continuous.
 - ii. T is continuous at the origin, in the sense that $x_n \to 0 \Rightarrow T(x_n) \to 0$;
 - iii. There exists a real number $k \ge 0$ with the property that $||T(x)|| \le k ||x||$ for every x in N;
 - iv. If $S = \{x : ||x|| \le 1\}$ is the closed unit sphere in N, then its image T(S) is a bounded set in N'.
 - b. Let B and B' be Banach spaces. If T is a continuous linear transformation of B onto B', then prove that image of each open sphere centered on the origin in B contains an open sphere centred on the origin in B'.

(OR)

Let B be a Banach space, and let M and N be closed linear subspaces of B such that $B = M \oplus N$. If z = x+y is the unique representation of a vector in B as a sum of vectors in M and N, then prove that the mapping P defined by P(z) = x is a projection on B whose range and null spaces are M and N respectively.

b. Let B be a Banach space and N a normed linear space if $\{T_n\}$ is a sequence in B (B,N) such that $T(x) = \lim_{n \to \infty} T_n(x)$ exists for each x in B then prove the T is a continuous linear transformation.

UNIT-IV

- 15. a. Let B be a complex Banach space whose norm obeys the parallelogram law, and if an inner product is defined on B by $4(x, y) = ||x + y||^2 ||x y||^2 + i||x + iy||^2 i||x iy||^2$, then prove that B is a Hilbert space.
 - b. If M is a closed linear subspace of a Hilbert space H, then prove that $H = M \oplus M^{\perp}$.

(OR)

16. a. Let H be a Hilbert space - prove that the adjoint operation $T \to T^*$ on B(H) has the following properties:

i.
$$(T_1 + T_2)^* = T_1^* + T_2^*$$
.

ii.
$$(\alpha T)^* = \overline{\alpha} T^*$$

iii.
$$(T_1T_2)^* = T_2 * T_1 *$$

iv.
$$T * * = T$$

$$v. \qquad ||T^*|| = ||T||$$

vi.
$$||T * T|| = ||T||^2$$
.

b. If T is an operator on a Hilbert space H, then prove that the following conditions are all equivalent to one another:

i.
$$T*T=I$$

ii.
$$(Tx,Ty) = (x,y)$$
 for all x,y;

iii.
$$||Tx|| = ||x||$$
 for all x.

PART II - FINAL

BRANCH: MATHEMATICS

Paper - III: OPERATIONS RESEARCH

(Revised Regulation from 2010-2011)

Max. Marks: 20

SECTION-A

Answer any Four questions. Each question carries 5 marks.

1. Define the term feasible solution and reduce the L.P.P. into its standard form:

Minimize
$$Z = -3x_1 + x_2 + x_3$$

$$x_1 - 2x_2 + x_3 \le 11$$
Subject to
$$-4x_1 + x_2 + 2x_3 \ge 3,$$

$$2x_1 - x_3 = -1; \ x_1, x_2 \ge 0$$

 x_3 unrestricted.

- 2. If $x_1 = 2$, $x_2 = 4$ and $x_3 = 1$ is a feasible solution of the system of equations $2x_1 x_2 + 2x_3 = 2$, $x_1 + 4x_2 = 18$ Reduce the given feasible solution to a basic feasible solution.
- 3. Write the dual of the L.P.P.

Minimize
$$Z = 4x_1 + 6x_2 + 18x_3$$

Subject to
$$x_1 + 3x_2 \ge 3$$
$$x_2 + 2x_3 \ge 5$$
$$x_1, x_2, x_3 \ge 0$$

4. Find an initial basic feasible solution of the transportation problem using the north west corner rule

				Supply
5	3	6	2	19
4	7	9	1	37
3	4	7	5	34
16	18	31	25	

- 5. Explain the formulation of Travelling salesman problem as Assignment problem.
- 6. Explain the fundamental EOQ problem.

Demand

- 7. Write two relations between average Queue length and average waiting time.
- 8. Discuss the probability distributions in Queueing systems.

SECTION - B

Answer one question from each unit. Each question carries 15 marks.

UNIT-I

9. a. Using graphical method find the maximum value of $Z = 10x_1 + 6x_2$

Subject to
$$5x_1 + 3x_2 \le 30$$

 $x_1 + 2x_2 \le 18$
 $x_1, x_2 \ge 0$

b. Use simplex method to solve

Maximize
$$Z = 3x_1 + 2x_2 + 5x_3$$

Subject to
$$x_1 + 2x_2 + x_3 \le 430$$

 $3x_1 + 2x_3 \le 460$
 $x_1 + 4x_3 \le 420$;
 $x_1, x_2, x_3 \ge 0$

(OR)

(2)

- 10. a. Show that dual of a dual is the primal.
 - b. State and prove fundamental theorem of Duality.

UNIT-II

11. Explain the least cost method and solve the transportation problem

				Supply
11	13	17	14	250
16	18	14	10	300
21	24	13	10	400
200	225	275	250	2-4
				(OD)

(OR)

- 12. a. Explain the vogel's approximation method.
 - b. Solve the assignment problem

Demand

10	25	15	20
15	30 .	5	15
35	20	12	24
17	25	24	20

UNIT-III

13. Use dynamic programming to solve the problem

Minimize
$$Z = y_1^2 + y_2^2 + y_3^2$$

Subject to
$$y_1 + y_2 + y_3 \ge 15$$
, $y_1, y_2, y_3 \ge 0$.

(OR)

14. If a project has the following time schedule given as

Activity	Time in weeks	Activity	Time in weeks
1-2	2	4-6	3
1-3	2	5-8	1
1-4	1	6-9	5
2-5	4	7-8	4
3-6	8	8-9	3
3-7	5	·	

Find the total float and critical path.

UNIT-IV

- 15. Write the operating characteristics of a Queueing system. a.
 - The rate of arrival of customers at a public telephone booth follow poisson distribution b. with an average time of 10 minutes between customer and the next. The duration of a phone call is exponential distribution with near time of 3 minutes then find
 - i. The probability that a person arriving at the booth will have to wait.
 - ii. Average length of non - empty queues.

(OR)

Explain the EOQ problem with finite Replenishment and write its characteristics.

PART II - Final

BRANCH: MATHEMATICS

Paper - IV: NUMBER THEORY

(Revised Regulation from 1999-2000)

Max. Marks: 20

SECTION - A

Answer any Four questions. Each questions carries 5 marks.

- 1. Prove that $\sum_{d/n} \phi(d) = n$.
- 2. State Euler Fermat theorem.
- 3. Define
 - i. Chebyshev's ψ function and
 - ii. Chebyshev's θ function.
- 4. Define character of a finite group G.
- 5. Let g be any real valued character mod k and let $A(n) = \sum_{d/n} y(d)$. Then prove that $A(n) \ge 0$ for all $n \ge 1$ and $A(n) \ge 1$ if n is a square.
- 6. Define
 - i. n is a quadratic residue modulo a prime p.
 - ii. n is note a quadratic residue modulo a prime p.
- 7. Let a and m be relativity prime integers. When do you say that a is a primitive root modulo m.

(1)

8. For a fixed $k \ge 1$, let $g(n) = \sum_{m=0}^{k-1} e^{\frac{2\pi i m^n}{k}}$. Then prove that $g(n) = \begin{cases} 0 & \text{if } k + n \\ k & \text{if } k / n \end{cases}$.

SECTION-B

Answer one question from each unit. Each question carries 15 marks.

UNIT - I

- 9. a. Let f be multiplicative function. Then, prove that f is completely multiplicative if and only if $f^{-1}(n) = \mu(n) \cdot f(n)$ for all $n \ge 1$.
 - b. State and prove Euler's summation formula.

(OR)

- 10. a. Assume that (a, m) = d. Then prove that the linear congruence $ax \equiv b \pmod{m}$ has solutions if and only if d/b.
 - b. State and prove chinese remainder theorem.

UNIT-II

11. For $n \ge L$, prove that $\frac{1}{6} \cdot \frac{n}{\log n} < \pi(n) < 6 \cdot \frac{n}{\log n}$.

(OR)

12. State and prove Selberg's asymptotic formula.

UNIT - III

13. Prove that a finite abelian group of order n - has exactly n - distinct characters.

(OR)

14. If the relation $\pi_a(x) \sim \frac{\pi(x)}{\phi(k)}$ as $x \to \infty$ holds for every integer 'a' relatively prime to k then prove that $\pi_a(x) \sim \pi_b(x)$ as $x \to \infty$ whenever (a,k) = (b,k) = 1.

UNIT-IV

15. Let $S_k(n) = \sum_{d'(n,k)} f(d)g\left(\frac{k}{d}\right)$ where f and g are multiplicative. Then prove that $S_{mk}(ab) = S_m(a).S_k(b)$ whenever (a,k)=(b,m)=1.

(OR)

16. a. Let (a,m) = 1. Then prove that a is a primitive root mod m if and only if the numbers $a, a^2, \dots, a^{\phi(m)}$ form a reduced residue system modulo m.

(2)

b. Let (a,m) = 1, let $f = \exp_m(a)$. Then prove that $\exp_m(a^k) = \frac{\exp_m(a)}{(k,f)}$.

PART - II: FINAL

BRANCH: MATHEMATICS

Paper -V - MATHEMATICAL STATISTICS

(Revised Regulation from 2010-2011)

Max. Marks: 20

SECTION-A

Answer any FOUR questions. Each questioncarries 5 marks.

- 1. Explain
 - i) Random variable.
 - ii) Stochastic Independence.
- 2. Let the random variable X have the pdf where $f(x) = \begin{cases} 2x & ; 0 < x < 1 \\ 0 & ; elsewhere \end{cases}$.

Find the Probability density function?

- 3. Illustrate Gamma and Chi-Square distribution
- 4. A box contains 6 red, 4 white and 5 black balls. A person draws 4 balls from the box at random. Find the probability that among the balls drawn there is at least one ball of each colour?
- 5. If X has the moment generating function $M(t) = e^{2t+32t^2}$ prove that X has a normal distribution with M=2 and $\sigma^2 = 64$.
- 6. Explain MLE of θ in Estimation.
- 7. Define
 - a) Testing of Hypothesis
 - b) Likelihood ratio test
- 8. Let $X_1, X_2, X_3, ... X_n$ be a random sample from U[0, θ] population. Obtain MVUE for θ .

SECTION-B

Answer ONE question form each Unit. Each question carries 15 marks.

UNIT-I

- 9. a) State and prove Chebyshev's Inequality.
 - b) Given the joint density function of X and Y as

$$f(x,y) = \begin{cases} \frac{1}{2}x \exp(-4) & ; & 0 < x < 2, y > 0 \\ 0 & ; & elsewhere \end{cases}$$

Find the distribution of X+Y?

(OR)

- 10. a) State and prove Baye's theorem.
 - b) Let X and Y have the Joint probability density function described as follows:

1 0/						
f(x,y)	2/15	4/15	3/15	1/15	1/15	4/15
(x,y)	(1,1)	(1,2)	(1,3)	(2,1)	(2,2)	(2,3)

and f(x, y) is equal to zero elsewhere. Find the Correlation Coefficient ρ ?

UNIT-II

11. Derive the Poisson distribution as a limiting case of Binomial distribution.

(OR)

12. Fit the Binomial distribution for the following data and Compare the theoretical frequency with actual one.

X	0	1	2	3	4	5
Frequency	2	14	20	34	22	8

UNIT-III

13. Explain Normal distribution and its properties.

(OR)

14. State and explain t and F distributions.

UNIT-IV

15. State and prove Neyman-Pearson Lemma

(OR)

16. Define Cramer-rao Inequality